


## WE CLAIM:

1. A compound having the structure of Formula I



FORMULA - I

its pharmaceutically acceptable salts, amides, enantiomers, diastereomers, N-oxides, prodrugs, metabolites or their polymorphs, wherein A is a straight or branched C<sub>1</sub>-C<sub>4</sub> alkyl chain; R is cinnamyl, benzyl, substituted benzyl, phenyl, mono- or disubstituted phenyl group substituted with the substituents independently selected from the group consisting of halogen, hydroxy, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>6</sub> alkoxy, trifluoromethyl, nitro, trifluoroalkoxy group, or (dihalodiphenyl) methyl,

2. The compound according to claim 1 wherein halogen is selected from the group consisting of chloro, fluoro, iodo; C<sub>1</sub>-C<sub>6</sub> alkyl is selected from the group consisting of methyl, ethyl, n-propyl, isopropyl, butyl, tert-butyl, and C<sub>1</sub>-C<sub>6</sub> alkoxy is selected from the group consisting of methoxy, ethoxy, n-propoxy, isopropoxy, and hexyloxy.


3. The compounds according to claim 1 selected from the group consisting of:

- 2-[3-{4-(2-Methoxyphenyl)piperazin-1-yl}propyl]-3a,4,7,7a-tetrahydro-1H-isoindole-1,3(2H)-dione (Compound 01).
- 2-[3-{4-(3-Chlorophenyl)piperazin-1-yl}propyl]-3a,4,7,7a-tetrahydro-1H-isoindole-1,3(2H)-dione (Compound 02).
- 2-[3-{4-(2-Methylphenyl)piperazin-1-yl}propyl]-3a,4,7,7a-tetrahydro-1H-isoindole-1,3(2H)-dione (Compound 03).

- Q1  
Cont
- 2-[3-{4-(4-Fluorophenyl)piperazin-1-yl}propyl]-3a,4,7,7a-tetrahydro-1H-isoindole-1,3(2H)-dione (Compound 04).
  - 2-[3-{4-(3-Trifluoromethylphenyl)piperazin-1-yl}propyl]-3a,4,7,7a-tetrahydro-1H-isoindole-1,3(2H)-dione (Compound 05).
  - 2-[3-{4-(2-Fluorophenyl)piperazin-1-yl}propyl]-3a,4,7,7a-tetrahydro-1H-isoindole-1,3(2H)-dione (Compound 06).
  - 2-[3-{4-(3,4-Dimethylphenyl)piperazin-1-yl}propyl]-3a,4,7,7a-tetrahydro-1H-isoindole-1,3(2H)-dione (Compound 07).
  - 2-[3-{4-(2-Methoxy-5-fluorophenyl)piperazin-1-yl}propyl]-3a,4,7,7a-tetrahydro-1H-isoindole-1,3(2H)-dione (Compound 08).
  - 2-[3-{4-(2-Ethylphenyl)piperazin-1-yl}propyl]-3a,4,7,7a-tetrahydro-1H-isoindole-1,3(2H)-dione (Compound 09).
  - 2-[3-{4-(2,4-Difluorophenyl)piperazin-1-yl}propyl]-3a,4,7,7a-tetrahydro-1H-isoindole-1,3(2H)-dione (Compound 10).
  - 2-[3-{4-(2-Ethoxyphenyl)piperazin-1-yl}propyl]-3a,4,7,7a-tetrahydro-1H-isoindole-1,3(2H)-dione (Compound 11).
  - 2-[3-{4-(2-Methyl-5-chlorophenyl)piperazin-1-yl}propyl]-3a,4,7,7a-tetrahydro-1H-isoindole-1,3(2H)-dione (Compound 12).
  - 2-[3-{4-(Phenyl)piperazin-1-yl}propyl]-3a,4,7,7a-tetrahydro-1H-isoindole-1,3(2H)-dione (Compound 13).
  - 2-[3-{4-(Benzyl)piperazin-1-yl}propyl]-3a,4,7,7a-tetrahydro-1H-isoindole-1,3(2H)-dione (Compound 14).
  - 2-[3-{4-(Cinnamyl)piperazin-1-yl}propyl]-3a,4,7,7a-tetrahydro-1H-isoindole-1,3(2H)-dione (Compound 15).
  - 2-[3-{4-(4-Nitrophenyl)piperazin-1-yl}propyl]-3a,4,7,7a-tetrahydro-1H-isoindole-1,3(2H)-dione (Compound 16).
  - 2-[3-{4-(3-Chloro-4-methylphenyl)piperazin-1-yl}propyl]-3a,4,7,7a-tetrahydro-1H-isoindole-1,3(2H)-dione (Compound 17).
  - 2-[3-{4-(4-Fluoro-2-methoxyphenyl)piperazin-1-yl}propyl]-3a,4,7,7a-tetrahydro-1H-isoindole-1,3(2H)-dione (Compound 18).
  - 2-[3-{4-(Bis-4-fluorophenyl)methylpiperazin-1-yl}propyl]-3a,4,7,7a-tetrahydro-1H-isoindole-1,3(2H)-dione (Compound 19).
  - 2-[3-{4-(2,4-Dichlorophenyl)piperazin-1-yl}propyl]-3a,4,7,7a-tetrahydro-1H-isoindole-1,3(2H)-dione (Compound 20).

- Q1  
con
- 2-[3-{4-(2,4-Dimethoxyphenyl)piperazin-1-yl}propyl]-3a,4,7,7a-tetrahydro-1H-isoindole-1,3(2H)-dione (Compound 21).
  - 2-[3-{4-(2,6-Dimethylphenyl)piperazin-1-yl}propyl]-3a,4,7,7a-tetrahydro-1H-isoindole-1,3(2H)-dione (Compound 22).
  - 2-[3-{4-(2-Isopropoxyphenyl)piperazin-1-yl}propyl]-3a,4,7,7a-tetrahydro-1H-isoindole-1,3(2H)-dione (Compound 23).
  - 2-[3-{4-(2-Propoxyphenyl)piperazin-1-yl}propyl]-3a,4,7,7a-tetrahydro-1H-isoindole-1,3(2H)-dione (Compound 24).
  - 2-[3-{4-(2-n-Hexyloxyphenyl)piperazin-1-yl}propyl]-3a,4,7,7a-tetrahydro-1H-isoindole-1,3(2H)-dione (Compound 25).
  - 2-[3-{4-(2,5-Dimethoxyphenyl)piperazin-1-yl}propyl]-3a,4,7,7a-tetrahydro-1H-isoindole-1,3(2H)-dione (Compound 26).
  - 2-[3-{4-(4-tert-Butylphenyl)piperazin-1-yl}propyl]-3a,4,7,7a-tetrahydro-1H-isoindole-1,3(2H)-dione (Compound 27).
  - 2-[3-{4-(2-Methoxy-6-hydroxyphenyl)piperazin-1-yl}propyl]-3a,4,7,7a-tetrahydro-1H-isoindole-1,3(2H)-dione (Compound 28).
  - 2-[3-{4-(2-Methoxyphenyl)piperazin-1-yl}-3-methylpropyl]-3a,4,7,7a-tetrahydro-1H-isoindole-1,3(2H)-dione (Compound 29).
  - 2-[3-{4-(2-Methoxyphenyl)piperazin-1-yl}-2-methylpropyl]-3a,4,7,7a-tetrahydro-1H-isoindole-1,3(2H)-dione (Compound 30).
  - 2-[3-{4-(2-Ethoxyphenyl)piperazin-1-yl}-3-methylpropyl]-3a,4,7,7a-tetrahydro-1H-isoindole-1,3(2H)-dione (Compound 31).


4. A method of selectively antagonizing  $\alpha_1$ -adrenergic receptors in a mammal comprising administering to said mammal a compound having the structure of Formula I



10  
FORMULA - I

its pharmaceutically acceptable salts, amides, enantiomers, diastereomers, N-oxides, prodrugs, metabolites or their polymorphs, wherein A is a straight or branched C<sub>1</sub>-C<sub>4</sub> alkyl chain; R is cinnamyl, benzyl, substituted benzyl, phenyl, mono- or disubstituted phenyl group substituted with the substituents independently selected from the group consisting of halogen, hydroxy, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>6</sub> alkoxy, trifluoromethyl, nitro, trifluoroalkoxy group, or (dihalodiphenyl) methyl.

5. A method for treating benign prostatic hyperplasia in a mammal comprising administering to said mammal a compound having the structure of Formula I



FORMULA - I


its pharmaceutically acceptable salts, amides, enantiomers, diastereomers, N-oxides, prodrugs, metabolites or their polymorphs, wherein A is a straight or branched C<sub>1</sub>-C<sub>4</sub> alkyl chain; R is cinnamyl, benzyl, substituted benzyl, phenyl, mono- or disubstituted with the substituents independently selected from the group consisting of halogen, hydroxy, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>6</sub> alkoxy, trifluoromethyl, nitro, trifluoroalkoxy group, or (dihalodiphenyl) methyl.

6. A pharmaceutical composition comprising the compound of claim 1 and a pharmaceutical acceptable carrier.

7. A method of selectively antagonizing  $\alpha_1$ -adrenergic receptors in a mammal comprising the step of administering to the said mammal the pharmaceutical composition according to claim 6.

8. A method for treating benign prostatic hyperplasia in a mammal comprising the step of administering to the said mammal the pharmaceutical composition according to claim 6.


9. A process for preparing a compound of Formula I



FORMULA - I

or its pharmaceutically acceptable salts, amides, enantiomers, diastereomers, N-oxides, prodrugs, metabolites or their polymorphs, wherein A is a straight or branched C<sub>1</sub>-C<sub>4</sub> alkyl chain; R is cinnamyl, benzyl, substituted benzyl, phenyl, mono- or disubstituted phenyl group substituted with the substituents independently selected from the group consisting of halogen, hydroxy, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>6</sub> alkoxy, trifluoromethyl, nitro, trifluoroalkoxy group, or (dihalodiphenyl) methyl, which comprises reacting a compound of Formula II, with piperazine derivatives of Formula III, as shown in Scheme I wherein A and R are the same as defined above.

10. A process for preparing a compound of Formula I



FORMULA - I

or its pharmaceutically acceptable salts, amides, enantiomers, diastereomers, N-oxides, prodrugs, metabolites or their polymorphs, wherein A is a straight or

a'  
cont

5 branched C<sub>1</sub>-C<sub>4</sub> alkyl chain; R is cinnamyl, benzyl, substituted benzyl, phenyl, mono- or disubstituted phenyl group substituted with the substituents independently selected from the group consisting of halogen, hydroxy, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>6</sub> alkoxy, trifluoromethyl, nitro, trifluoroalkoxy group or (dihalodiphenyl) methyl, which comprises reacting 1-(ω-haloalkyl)cis-3a,4,7,7a-tetrahydrophthalimide of Formula IV, wherein A is the same as defined above, with a compound of Formula V, wherein R is the same as defined above, as shown in Scheme II.